If it is desired to modify the classification of Ball, *Gehringia* should be transferred to the Subfamily Carabinae, of which it may form a Tribe, Gehringiini. (The Subfamily Carabinae of Ball is approximately equivalent to the Anisochaeta of Jeannel.) The removal of *Gehringia* would leave the Isochaeta as a relatively homogeneous group, with the structure of the tibial spurs, antenna-cleaner, and the chaetotaxy of the labrum as common characters. I believe that these characters imply a real relationship, and that the Subfamilies Paussinae, Metriinae, and Trachypachydinae (excluding Gehringiini) of Ball should be united to form a Subfamily Paussinae, approximately equivalent to Jeannel's Isochaeta.

LITERATURE CITED

BALL, G. E.

1960. Carabidae, pp. 55-181, in R. H. Arnett, The Beetles of the United States. Catholic University Press, 1112 pp., Washington, D. C.

DARLINGTON, P. J.

1933. A new tribe of Carabidae from western United States. Pan-Pacific Ent. IX (3), 110-114.

JEANNEL, R.

1941. Coléoptères carabiques (1) Faune de France, 39:1-571, Paris.

LENG, C. W.

1920. Catalogue of the Coleoptera of America, North of Mexico. 470 pp., J. D. Sherman, Jr., Mount Vernon, N. Y.

LINDROTH, C. H.

1960. The ground beetles of Canada and Alaska (2), Opusc. Ent. Suppl. XX:1-200, Lund, Sweden.

Tipe Marian

CEUTORHYNCHUS ASSIMILIS (PAYK.), NEW TO EASTERN NORTH AMERICA (COLEOPTERA: CURCULIONIDAE).

The finding of C. assimilis (Payk.) May 16, 1960, on turnips, in Henderson Co., North Carolina, represents the first record of this species in eastern North America. A serious pest of Cruciferae, the species was previously found only in the Pacific Northwest, in Washington, Oregon, British Columbia, and northern California. Rose ELLA WARNER, Ent. Res. Div., A.R.S., U. S. Department of Agriculture, Washington, D. C.